## Rabbit CD40 Ligand / TNFSF5 Protein, His, Flag Tag (active trimer) (MALS verified)

Catalog # CD0-R52D9



## Synonym

CD40LG,CD154,CD40L,HIGM1,IGM,IMD3,T-BAM,TNFSF5,TRAP,gp39

### **Source**

Rabbit CD40 Ligand, His,Flag Tag(CD0-R52D9) is expressed from human 293 cells (HEK293). It contains AA Gly 116 - Leu 261 (Accession # G1SKP7-1). Predicted N-terminus: His

### **Molecular Characterization**

This protein carries a polyhistidine tag at the N-terminus, followed by a flag tag.

The protein has a calculated MW of 49.9 kDa. The protein migrates as 50-55 kDa under non-reducing (NR) condition (SDS-PAGE) due to glycosylation.

#### **Endotoxin**

Less than 1.0 EU per µg by the LAL method.

## **Purity**

>90% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

#### **Formulation**

Lyophilized from  $0.22~\mu m$  filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

#### Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

## Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.


## **SDS-PAGE**



Rabbit CD40 Ligand, His,Flag Tag on SDS-PAGE under non-reducing (NR) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%.

## **Bioactivity-ELISA**

#### **SEC-MALS**




The purity of Rabbit CD40 Ligand, His,Flag Tag (Cat. No. CD0-R52D9) is more than 90% and the molecular weight of this protein is around 45-60 kDa verified by SEC-MALS.

Report

## Rabbit CD40 Ligand / TNFSF5 Protein, His,Flag Tag (active trimer) (MALS verified)

Catalog # CD0-R52D9





Rabbit CD40 Ligand, His,Flag Tag ELISA

0.5 µg of Human CD40 Protein, Fc Tag (MALS verified) per well

2


EC50=7.04 ng/mL


0.1 1 10 100 1000

Rabbit CD40 Ligand, His,Flag Tag Conc. (ng/mL)

Immobilized Human CD40, Mouse IgG2a Fc Tag, low endotoxin (MALS verified) (Cat. No. CD0-H525a) at 5 μg/mL (100 μL/well) can bind Rabbit CD40 Ligand, His,Flag Tag (Cat. No. CD0-R52D9) with a linear range of 1-39 ng/mL (QC tested).

Immobilized Human CD40 Protein, Fc Tag (MALS verified) (Cat. No. CD0-H5253) at 5  $\mu$ g/mL (100  $\mu$ L/well) can bind Rabbit CD40 Ligand, His,Flag Tag (Cat. No. CD0-R52D9) with a linear range of 0.6-39 ng/mL (Routinely tested).





Immobilized Rabbit CD40 Ligand, His,Flag Tag (Cat. No. CD0-R52D9) at 5  $\mu$ g/mL (100  $\mu$ L/well) can bind Human CD40 Protein, Fc Tag (MALS verified) (Cat. No. CD0-H5253) with a linear range of 0.156-20  $\mu$ g/mL (Routinely tested).

Immobilized Rabbit CD40, His Tag (Cat. No. CD0-R52H9) at 5  $\mu$ g/mL (100  $\mu$ L/well) can bind Rabbit CD40 Ligand, His,Flag Tag (Cat. No. CD0-R52D9) with a linear range of 1-5 ng/mL (Routinely tested).

## **Background**

CD40 ligand is also known as CD40L, CD154, TNFSF5 and T-cell antigen Gp39, is a single-pass type I I membrane protein which belongs to the TNF superfamily of molecules. CD40 ligand is expressed predominantly on activated CD4+ T lymphocytes, and also found in other types of cells, including platelets, mast cells, macrophages, basophils, NK cells, B lymphocytes, as well as non-haematopoietic cells (smooth muscle cells, endothelial cells, and epithelial cells). Although all monomeric, dimeric and trimeric forms of soluble CD40 ligand can bind to CD40, the trimeric form of soluble CD40 ligand has the most potent biological activity through oligomerization of cell surface CD40, a common feature of TNF receptor family members.

CD40 ligand binds to CD40 on antigen-presenting cells (APC), which leads to many effects depending on the target cell type. In general, CD40 ligand plays the role of a costimulatory molecule and induces activation in APC in association with T cell receptor stimulation by MHC molecules on the APC. In total CD40 ligand has three binding partners: CD40, α5β1 integrin and αIIbβ3. CD40 ligand regulates B cell function by engaging CD40 on the B cell surface. A defect in this gene results in an inability to undergo immunoglobulin class switch and is associated with hyper IgM syndrome.

# **Clinical and Translational Updates**

