



Synonym

CD4,CD4mut,LEU3

Source

Human CD4 Protein, Fc Tag(CD4-H5259) is expressed from human 293 cells (HEK293). It contains AA Lys 26 - Pro 396 (Accession # [AAH25782](#)).

Predicted N-terminus: Lys 26

Molecular Characterization

CD4(Lys 26 - Pro 396) AAH25782	Fc(Pro 100 - Lys 330) P01857
-----------------------------------	---------------------------------

This protein carries a human IgG1 Fc tag at the C-terminus.

The protein has a calculated MW of 67.5 kDa. The protein migrates as 70-80 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 0.2 EU per µg by the LAL method.

Purity

>95% as determined by SDS-PAGE.

>95% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 µm filtered solution in Tris, Glycine and NaCl, pH7.5 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 12 months under sterile conditions after reconstitution.


SDS-PAGE



Human CD4 Protein, Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

Bioactivity-ELISA

SEC-MALS



The purity of Human CD4 Protein, Fc Tag (Cat. No. CD4-H5259) is more than 95% and the molecular weight of this protein is around 150-183 kDa verified by SEC-MALS.

[Report](#)

Discounts, Gifts,
and more!




Human CD4 Protein, Fc Tag (MALS verified)


Catalog # CD4-H5259



BIOSYSTEMS
Acro



Immobilized Human CD4 Protein, Fc Tag (Cat. No. CD4-H5259) at 1 µg/mL (100 µL/well) can bind Ibalizumab (Human IgG4) with a linear range of 0.2-3 ng/mL (QC tested).



Immobilized HIV-1 [HIV-1/Clade B/C (CN54)] GP120, His Tag (Cat. No. GP4-V15227) at 2 µg/mL (100 µL/well) can bind Human CD4 Protein, Fc Tag (Cat. No. CD4-H5259) with a linear range of 1-31 ng/mL (Routinely tested).

Background

T-cell surface glycoprotein CD4 is also known as T-cell surface antigen T4/Leu-3. CD4 contains three Ig-like C2-type (immunoglobulin-like) domains and one Ig-like V-type (immunoglobulin-like) domain. CD4 is accessory protein for MHC class-II antigen/T-cell receptor interaction. CD4 induces the aggregation of lipid rafts. CD4 is a primary receptor used by HIV-1 to gain entry into host T cells. HIV infection leads to a progressive reduction of the number of T cells possessing CD4 receptors. Therefore, medical professionals refer to the CD4 count to decide when to begin treatment for HIV-infected patients.

Clinical and Translational Updates

Discounts, Gifts,
and more!

