

Synonym

CEACAM-5,CD66e,CEA,Meconium antigen 100

Source

Mouse CEACAM-5, Fc Tag(CE5-M5253) is expressed from human 293 cells (HEK293). It contains AA Gln 35 - Glu 947 (Accession # Q3UKK2-1).

Molecular Characterization

CEACAM-5(Gln 35 - Glu 947) Fc(Pro 100 - Lys 330)
Q3UKK2-1 P01857

This protein carries a human IgG1 Fc tag at the C-terminus.

The protein has a calculated MW of 129.2 kDa. The protein migrates as 135-180 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per μg by the LAL method.

Purity

>90% as determined by SDS-PAGE.

Formulation

Lyophilized from $0.22~\mu m$ filtered solution in 50~mM Tris, 100~mM Glycine, 25~mM Arginine, 150~mM NaCl, pH7.5 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:


- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE



Mouse CEACAM-5, Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%.

SEC-MALS



The purity of Mouse CEACAM-5, Fc Tag (Cat. No. CE5-M5253) is more than 85% and the molecular weight of this protein is around 280-360 kDa verified by SEC-MALS.


Report

Background

Carcinoembryonic antigen-related cell adhesion molecule 5 (CEACAM5) is also Carcinoembryonic antigen (CEA), Meconium antigen 100, CD antigen CD66e, CEACAM5 belongs to the immunoglobulin superfamily and CEA family. CEACAM5 contains seven Ig-like (immunoglobulin-like) domains. CEACAM5 is



Mouse CEACAM-5 Protein, Fc Tag (MALS verified)





homodimer protein which binding of E.coli Dr adhesins leads to dissociation of the homodimer. CEACAM5 is cell surface glycoprotein that plays a role in cell adhesion and in intracellular signaling. CEACAM5 is receptor for E.coli Dr adhesins.

Clinical and Translational Updates

