

Mouse CTLA-4 / CD152 Protein, His Tag (MALS verified)

Catalog # CT4-M52H5



BIOSYSTEMS
Acro
Surprise Inside!

Synonym

CTLA4,CD152

Source

Mouse CTLA-4, His Tag(CT4-M52H5) is expressed from human 293 cells (HEK293). It contains AA Glu 36 - Phe 162 (Accession # [NP_033973](#)).

Predicted N-terminus: Glu 36

Molecular Characterization

CTLA-4(Glu 36 - Phe 162)
NP_033973 Poly-his

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 15.9 kDa. The protein migrates as 23-30 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>95% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.


SDS-PAGE



Mouse CTLA-4, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95% (With [Star Ribbon Pre-stained Protein Marker](#)).

Bioactivity-ELISA

SEC-MALS



The purity of Mouse CTLA-4, His Tag (Cat. No. CT4-M52H5) is more than 90% and the molecular weight of this protein is around 40-50 kDa verified by SEC-MALS.

[Report](#)

Discounts, Gifts,
and more!




➤ www.acrobiosystems.com


9/19/2024

Mouse CTLA-4 / CD152 Protein, His Tag (MALS verified)

Catalog # CT4-M52H5



Immobilized Mouse B7-1, Fc Tag (Cat. No. CD0-M5259) at 2 µg/mL (100 µL/well) can bind Mouse CTLA-4, His Tag (Cat. No. CT4-M52H5) with a linear range of 0.1-2 ng/mL (QC tested).



Immobilized Mouse B7-2, Fc Tag (Cat. No. CD6-M5251) at 5 µg/mL (100 µL/well) can bind Mouse CTLA-4, His Tag (Cat. No. CT4-M52H5) with a linear range of 0.4-13 ng/mL (Routinely tested).

Background

CTLA-4 (Cytotoxic T-Lymphocyte Antigen 4) is also known as CD152 (Cluster of differentiation 152), is a protein receptor that downregulates the immune system. CTLA4 is a member of the immunoglobulin superfamily, which is expressed on the surface of Helper T cells and transmits an inhibitory signal to T cells. The protein contains an extracellular V domain, a transmembrane domain, and a cytoplasmic tail. Alternate splice variants, encoding different isoforms. CTLA4 is similar to the T-cell co-stimulatory protein, CD28, and both molecules bind to CD80 and CD86, also called B7-1 and B7-2 respectively, on antigen-presenting cells. CTLA4 transmits an inhibitory signal to T cells, whereas CD28 transmits a stimulatory signal. Intracellular CTLA4 is also found in regulatory T cells and may be important to their function. Fusion proteins of CTLA4 and antibodies (CTLA4-Ig) have been used in clinical trials for rheumatoid arthritis.

Clinical and Translational Updates

Discounts, Gifts,
and more!

