

## **Synonym**

Delta3,delta-like 3 (Drosophila),delta-like protein 3,DLL3,Pudgy,SCDO1,SCDO1delta3

#### Source

Rat DLL3, His Tag(DL3-R52H3) is expressed from human 293 cells (HEK293). It contains AA Ala 25 - Ala 488 (Accession # <u>088671-1</u>).

Predicted N-terminus: Ala 25

### **Molecular Characterization**

DLL3(Ala 25 - Ala 488) O88671-1

Poly-his

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 50.1 kDa. The protein migrates as 50-60 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

#### **Endotoxin**

Less than 1.0 EU per µg by the LAL method.

# **Purity**

>90% as determined by SDS-PAGE.

## **Formulation**

Lyophilized from 0.22  $\mu m$  filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

#### Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

### Storage


For long term storage, the product should be stored at lyophilized state at -20 $^{\circ}$ C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

## **SDS-PAGE**



Rat DLL3, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%.


## **Bioactivity-ELISA**



# Rat DLL3 Protein, His Tag

Catalog # DL3-R52H3





Anti-DLL3 Antibody (specific Binding DSL of DLL3), Human IgG1 Conc. (ng/mL)

Immobilized Rat DLL3, His Tag (Cat. No. DL3-R52H3) at 1  $\mu$ g/mL (100  $\mu$ L/well) can bind Anti-DLL3 Antibody (specific Binding DSL of DLL3), Human IgG1 with a linear range of 0.1-4 ng/mL (QC tested).

# **Background**

Delta-like protein 3 (DLL3) is a transmembrane protein that belongs to the Delta/Serrate/Lag-2 (DSL) family of Notch ligands. May be required to divert neurons along a specific differentiation pathway. Plays a role in the formation of somite boundaries during segmentation of the paraxial mesoderm. DLL3 protein is expressed on the surface of tumor cells but not in normal adult tissues.

## **Clinical and Translational Updates**

Please contact us via <u>TechSupport@acrobiosystems.com</u> if you have any question on this product.

