Catalog # DNP-FM487



Source

Mouse IgG2a Antibody Isotype Control is a monoclonal antibody recombinantly expressed from human 293 cells (HEK293), which provides higher batch consistency and long term security of supply.

Application

Flow Cytometry.

Species

Mouse

Isotype

Mouse IgG2a | Mouse kappa

Specificity

Specifically reacts with DNP (Dinitrophenyl) and DNP conjugated proteins, which is normally not present in Humans or animals. Therefore, DNP-FM487 does not react with any antigen of Human or animal cells.

Immunogen

DNP (Dinitrophenyl).

Conjugate


FITC

Excitation source: 488 nm spectral line, argon-ion laser

Excitation Wavelength: 488 nm

Emission Wavelength: 535 nm

Bioactivity-FACS



Recommended Dilution

1:50

Formulation

Lyophilized from 0.22 μ m filtered solution in PBS, 0.03% Proclin300, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.


Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please protect from light and avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 24 months in lyophilized state;
- -70°C for 12 months after reconstitution;
- 2-8°C for 12 months after reconstitution.



5e5 of PBMCs were stained with FITC-Labeled Mouse IgG2a Antibody Isotype Control (Cat. No. DNP-FM487), washed and then analyzed with





Catalog # DNP-FM487



FACS. FITC signal was used to evaluate the non-specific binding activity to PBMCs (QC tested).

Background

A hapten is a small molecule that can elicit an immune response only when conjugated with a large carrier such as a protein. Typical haptens include drugs, urushiol, quinone, steroids, etc. Peptides and non-protein antigens usually need conjugating to a carrier protein (such as BSA (bovine serum albumin) or KLH (keyhole limpet hemocyanin) to become good immunogens). Additionally, haptens should be administered with an adjuvant to ensure a high quality immune response. It is important that the hapten design (preserving greatly the chemical structure and spatial conformation of target compound), selection of the appropriate carrier protein and the conjugation method are key conditions for the desired specificity anti-hapten antibodies. We design anti-hapten antibodies based on the HaptenDB information.

Clinical and Translational Updates



>>> www.acrobiosystems.com

