



Synonym

IL7Ra,CD127

Source

Human IL-7 R alpha, Fc Tag(ILA-H525a) is expressed from human 293 cells (HEK293). It contains AA Glu 21 - Asp 239 (Accession # [P16871-1](#)).

Molecular Characterization

IL-7 R alpha(Glu 21 - Asp 239) P16871-1	Fc(Pro 100 - Lys 330) P01857
--	---------------------------------

This protein carries a human IgG1 Fc tag at the C-terminus.

The protein has a calculated MW of 51.8 kDa. The protein migrates as 60-70 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>95% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE




Human IL-7 R alpha, Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

Bioactivity-ELISA

Discounts, Gifts,
and more!





Immobilized Human IL-7 Protein at 2 µg/mL (100 µL/well) can bind Human IL-7 R alpha, Fc Tag (Cat. No. ILA-H525a) with a linear range of 0.004-0.125 µg/mL (QC tested).

Clinical and Translational Updates

Discounts, Gifts,
and more!

