Human Integrin alpha V beta 6 (ITGAV&ITGB6) Heterodimer Protein, His Tag&Tag Free (MALS verified)

Catalog # IT6-H52E1





Synonym

Integrin alpha V beta 6,ITGAV&ITGB6

Source

Human ITGAV&ITGB6 Heterodimer Protein, His Tag&Tag Free(IT6-H52E1) is expressed from human 293 cells (HEK293). It contains AA Phe 31 - Val 992 (ITGAV) & Gly 22 - Asn 707 (ITGB6) (Accession # NP 002201.1 (ITGAV) & AAI21179.1 (ITGB6)).

Predicted N-terminus: Phe 31 (ITGAV) & Gly 22 (ITGB6)

Molecular Characterization

ITGAV (Phe 31 - Val 992) NP_002201.1	Acidic Tail	Poly-his
ITGB6 (Gly 22 - Asn 707) AAI21179.1	Basic Tail	

Human ITGAV&ITGB6 Heterodimer Protein, His Tag&Tag Free, produced by co-expression of ITGAV and ITGB6, has a calculated MW of 112.9 kDa (ITGAV) & 79.6 kDa (ITGB6). Subunit ITGAV is fused with an acidic tail at the C-terminus and followed by a polyhistidine tag and subunit ITGB6 contains no tag but a basic tail at the C-terminus. The protein migrates as 85 kDa (ITGB6) and 140 kDa (ITGAV) when calibrated against Star Ribbon Pre-stained Protein Marker under non-reducing (NR) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per μg by the LAL method.

Purity

>95% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in 50 mM Tris, 150 mM NaCl, pH7.5 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.


SDS-PAGE



Human ITGAV&ITGB6 Heterodimer Protein, His Tag&Tag Free on SDS-PAGE under non-reducing (NR) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95% (With Star Ribbon Pre-stained Protein Marker).

Bioactivity-ELISA

SEC-MALS



The purity of Human ITGAV&ITGB6 Heterodimer Protein, His Tag&Tag Free (Cat. No. IT6-H52E1) is more than 85% and the molecular weight of this protein is around 200-240 kDa verified by SEC-MALS. Report




Human Integrin alpha V beta 6 (ITGAV&ITGB6) Heterodimer Protein, His Tag&Tag Free (MALS verified)

Catalog # IT6-H52E1




Human ITGAV&ITGB6 Heterodimer Protein, His Tag&Tag Free ELISA 0.5 μ g of Human ITGAV&ITGB6 Heterodimer Protein, His Tag&Tag Free per well



Biotinylated Human Latent TGF-Beta 1, His, Avitag Conc. (ng/mL)

Immobilized Human ITGAV&ITGB6 Heterodimer Protein, His Tag&Tag Free (Cat. No. IT6-H52E1) at 5 μ g/mL (100 μ L/well) can bind Biotinylated Human Latent TGF-Beta 1, His,Avitag (Cat. No. TG1-H82Qb) with a linear range of 0.5-4 ng/mL (QC tested).


Human ITGAV&ITGB6 Heterodimer Protein, His Tag&Tag Free ELISA $1~\mu g$ of Human Fibronectin per well



Human ITGAV&ITGB6 Heterodimer Protein, His Tag&Tag Free Conc. (μg/mL)

Immobilized Human Fibronectin at 10 μ g/mL (100 μ L/well) can bind Human ITGAV&ITGB6 Heterodimer Protein, His Tag&Tag Free (Cat. No. IT6-H52E1) with a linear range of 0.313-2.5 μ g/mL (Routinely tested).

Bioactivity-SPR



Fibronectin fragment, premium grade (Cat. No. FIN-H5113) immobilized on CM5 Chip can bind Human ITGAV&ITGB6 Heterodimer Protein, His Tag&Tag Free (Cat. No. IT6-H52E1) with an affinity constant of 1.23 μ M as determined in a SPR assay (Biacore 8K) (Routinely tested).

Background

Integrin alpha V beta 6 is a heterodimer of beta-6 associating with alpha-V. Integrin alpha-V beta-6 is a receptor for fibronectin and cytotactin. It recognizes the sequence R-G-D in its ligands. Internalisation of integrin alpha-V beta-6 via clathrin-mediated endocytosis promotes carcinoma cell invasion. Also, Integrin alpha-V beta-6 acts as a receptor for coxsackievirus A9 and coxsackievirus B1 as well as herpes simplex virus-1/HHV-1. Furthermore, it binds the TGF-beta latency-associated peptide (LAP) and activates TGF-beta 1 or TGF-beta 3 from large latent complexes. This activation requires interaction with LTBP-1 and fibronectin, and is enhanced by PAR-1.

Clinical and Translational Updates

