



Synonym

LAG3,CD223,FDC

Source

Biotinylated Human LAG-3, Mouse IgG2a Fc,Avitag(LA3-H82F3) is expressed from human 293 cells (HEK293). It contains AA Leu 23 - Leu 450 (Accession # [P18627-1](#)).

Predicted N-terminus: Leu 23

Molecular Characterization

LAG-3(Leu 23 - Leu 450) P18627-1	mFc(Glu 98 - Lys 330) P01863	Avi
-------------------------------------	---------------------------------	-----

This protein carries a mouse IgG1 Fc tag at the C-terminus.

The protein has a calculated MW of 75.4 kDa. The protein migrates as 100 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Labeling

Biotinylation of this product is performed using Avitag™ technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.

Protein Ratio

Passed as determined by the HABA assay / binding ELISA.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>95% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 µm filtered solution in Tris with Glycine, Arginine and NaCl, pH7.5 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.


SDS-PAGE



Biotinylated Human LAG-3, Mouse IgG2a Fc,Avitag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

Bioactivity-ELISA

SEC-MALS




The purity of Biotinylated Human LAG-3, Mouse IgG2a Fc,Avitag (Cat. No. LA3-H82F3) is more than 90% and the molecular weight of this protein is around 170-200 kDa verified by SEC-MALS.

[Report](#)


Discounts, Gifts,
and more!





Immobilized Madarex LAG-3 MAb, Human IgG1 at 2 µg/mL (100 µL/well) can bind Biotinylated Human LAG-3, Mouse IgG2a Fc,Avitag (Cat. No. LA3-H82F3) with a linear range of 0.4-13 ng/mL (QC tested).

Bioactivity-FACS



Flow Cytometry assay shows that Biotinylated Human LAG-3, Mouse IgG2a Fc,Avitag (Cat. No. LA3-H82F3) can bind to daudi cell surface. The concentration of LAG-3 used is 0.3 µg/mL (Routinely tested).

Background

Lymphocyte activation gene 3 protein (LAG3) is also known as CD antigen CD223 and protein FDC, which belongs to immunoglobulin (Ig) superfamily and contains 4 extracellular Ig-like domains. The LAG3 gene contains 8 exons. The sequence data, exon/intron organization, and chromosomal localization all indicate a close relationship of LAG3 to CD4. LAG3 /CD223 involved in lymphocyte activation. LAG3 /CD223 binds to HLA class-II antigens.

Clinical and Translational Updates

Please contact us via TechSupport@acrobiosystems.com if you have any question on this product.

Discounts, Gifts,
and more!

