

Synonym

SLC39A6,LIV-1,ZIP6,Zinc transporter ZIP6,ZIP-6

Source

Rat LIV-1, His Tag (LV1-R52H9) is expressed from human 293 cells (HEK293).

It contains AA Leu 21 - Trp 311 (Accession # [Q4V887-1](#)).

Predicted N-terminus: Leu 21

Molecular Characterization

LIV1(Leu 21 - Trp 311)
Q4V887-1 Poly-his

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 34.6 kDa. The protein migrates as 55-65 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>90% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4. Normally trehalose is added as protectant before lyophilization.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.


Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE

Rat LIV-1, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 90%.

Background

LIV-1 is also known as SLC39A6, ZIP-6 and Zinc transporter ZIP6. May act as a zinc-influx transporter. Highly expressed in the breast, prostate, placenta, kidney, pituitary and corpus callosum. Weakly expressed in heart and intestine. Also highly expressed in cells derived from an adenocarcinoma of the cervix and lung carcinoma. Up-regulated by estrogen in breast cancer cells lines.

Clinical and Translational Updates

Please contact us via TechSupport@acrobiosystems.com if you have any question on this product.