Catalog # MCF-H5247



#### Synonym

M-CSF,CSF-1,Lanimostim

#### Source

Human M-CSF, His Tag(MCF-H5247) is expressed from human 293 cells (HEK293). It contains AA Glu 33 - Arg 255 (Accession # <u>P09603-1</u>). Predicted N-terminus: His

## **Molecular Characterization**

M-CSF(Glu 33 - Arg 255) Poly-his P09603-1

This protein carries a polyhistidine tag at the N-terminus.

The protein has a calculated MW of 26.0 kDa. The protein migrates as 36-45 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

### Endotoxin

Less than 1.0 EU per  $\mu$ g by the LAL method.

# Purity

>95% as determined by SDS-PAGE.

#### Formulation

Lyophilized from 0.22  $\mu$ m filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

#### Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

#### Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- $70^{\circ}$ C for 3 months under sterile conditions after reconstitution.

# **SDS-PAGE**



Human M-CSF, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

#### **Bioactivity-ELISA**




>>> www.acrobiosystems.com




# Human M-CSF / CSF-1 Protein, His Tag

Catalog # MCF-H5247




Immobilized Human M-CSF, His Tag (Cat. No. MCF-H5247) at 5  $\mu$ g/mL (100  $\mu$ L/well) can bind Human M-CSF R, Fc Tag (Cat. No. CSR-H5258) with a linear range of 0.1-4 ng/mL (QC tested).




Immobilized Human M-CSF, His Tag (Cat. No. MCF-H5247) at 5  $\mu$ g/mL (100  $\mu$ L/well) can bind Cynomolgus M-CSF R, Fc Tag (Cat. No. CSR-C5252) with a linear range of 1-78 ng/mL (Routinely tested).


# Bioactivity-SPR



Human M-CSF, His Tag (Cat. No. MCF-H5247) captured on CM5 Chip via anti-His antibody can bind Human M-CSF R, Fc Tag, low endotoxin (Cat. No. CSR-H5258) with an affinity constant of 1.9 nM as determined in SPR assay (Biacore T200) (Routinely tested).

#### **Bioactivity-BLI**







Loaded Human M-CSF, His Tag (Cat. No. MCF-H5247) on HIS1K Biosensor, can bind Human M-CSF R, Fc Tag, low endotoxin (Cat. No. CSR-H5258) with

Loaded Human M-CSF R, Fc Tag, low endotoxin (Cat. No. CSR-H5258) on Protein A Biosensor, can bind Human M-CSF, His Tag (Cat. No. MCF-H5247)






# Human M-CSF / CSF-1 Protein, His Tag

Catalog # MCF-H5247


an affinity constant of 3.29 nM as determined in BLI assay (ForteBio Octet Red96e) (Routinely tested).


#### **Bioactivity-Bioactivity CELL BASE**





Human M-CSF, His Tag (Cat. No. MCF-H5247) stimulates proliferation of RAW264.7 cells. The EC50 for this effect is 4.40-5.27 ng/mL (Routinely tested).





The bio-activity of Human M-CSF / CSF-1 Protein, His Tag (Cat. No. MCF-H5247) was determined by dose-dependent stimulation of the proliferation of M-NFS-60 cells. The EC50 for this effect is 7.747 ng/mL (Routinely tested).

#### Background

The colony stimulating factor 1 (CSF1), also known as macrophage colony-stimulating factor (M-CSF), is a secreted cytokine which influences hematopoietic stem cells to differentiate into macrophages or other related cell types. Eukaryotic cells also produce M-CSF in order to combat intercellular viral infection. It is one of the three experimentally described colony-stimulating factors. M-CSF binds to the colony stimulating factor 1 receptor. Macrophage colony-stimulating factor has been shown to interact with PIK3R2. M-CSF (or CSF-1) is a hematopoietic growth factor that is involved in the proliferation, differentiation, and survival of monocytes, macrophages, and bone marrow progenitor cells. Locally produced M-CSF in the vessel wall contributes to the development and progression of atherosclerosis.

Luminescence(RLU)

#### **Clinical and Translational Updates**

Please contact us via <u>TechSupport@acrobiosystems.com</u> if you have any question on this product.



with an affinity constant of 0.365 nM as determined in BLI assay (ForteBio Octet Red96e) (Routinely tested).



>>> www.acrobiosystems.com

