



Synonym

Mucin 1, MUC1, CD227, EMA, H23AG, KL-6, MAM6, MUC-1, SEC, MUC-1, X, MUC1, ZD, PEM, PEMT, PUM, CA15-3, Episialin

Source

Human Mucin-1 (1098-1158), Mouse IgG2a Fc Tag (MU1-H5259) is expressed from human 293 cells (HEK293). It contains AA Ser 1098 - Gly 1158

(Accession # [P15941-1](#)).

Predicted N-terminus: Ser 1098

Molecular Characterization

Mucin-1(Ser 1098 - Gly 1158) P15941-1	mFc(Glu 98 - Lys 330) P01863
--	---------------------------------

This protein carries a mouse IgG2a Fc tag at the C-terminus.

The protein has a calculated MW of 33.4 kDa. The protein migrates as 43-53 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 0.1 EU per µg by the LAL method.

Purity

>95% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in 50 mM Tris, 100 mM Glycine, 25 mM Arginine, 150 mM NaCl, pH7.5 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE



Human Mucin-1 (1098-1158), Mouse IgG2a Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95% (With [Star Ribbon Pre-stained Protein Marker](#)).

Background

Membrane mucins have several functions in epithelial cells including cytoprotection, extravasation during metastases, maintenance of luminal structure, and signal transduction. MUC17, contains an extended, repetitive extracellular glycosylation domain and a carboxyl terminus with two EGF-like domains, a SEA module

Discounts, Gifts,
and more!



Human Mucin-1 / MUC-1 (1098-1158) Protein, Mouse IgG2a Fc Tag

Catalog # MU1-H5259



BIOSYSTEMS
Acro

domain, a transmembrane domain, and a cytoplasmic domain with potential serine and tyrosine phosphorylation sites. Interacts via its C-terminus with PDZK1 and this interaction appears important for proper localization. Probably plays a role in maintaining homeostasis on mucosal surfaces.

Clinical and Translational Updates

Discounts, Gifts,
and more!



➤ www.acrobiosystems.com

7/24/2024