

**Synonym**

PDCD1,PD1,CD279,SLEB2

**Source**

Human PD-1 Protein, His Tag(PD1-H522a) is expressed from human 293 cells (HEK293). It contains AA Leu 25 - Gln 167 (Accession # [NP\\_005009.2](#)).

Predicted N-terminus: Leu 25

**Molecular Characterization**

PD-1(Leu 25 - Gln 167)  
NP\_005009.2 Poly-his

This protein carries a polyhistidine tag at the C-terminus, and has a calculated MW of 16.8 kDa. The N-terminus Sequence Analysis is Leu 25. The reducing (R) protein migrates as 31-44 kDa in SDS-PAGE due to glycosylation.

**Endotoxin**

Less than 0.01 EU per µg by the LAL method.

**Purity**

>95% as determined by SDS-PAGE.

**Formulation**

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

**Reconstitution**

Please see Certificate of Analysis for specific instructions.

*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.*


**Storage**

For long term storage, the product should be stored at lyophilized state at -20°C or lower.


*Please avoid repeated freeze-thaw cycles.*

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.


**SDS-PAGE**

Human PD-1 Protein, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

**Bioactivity-ELISA****SEC-MALS**


The purity of Human PD-1 Protein, His Tag (Cat. No. PD1-H522a) is more than 85% and the molecular weight of this protein is around 25-40 kDa verified by SEC-MALS.

[Report](#)




Immobilized Human PD-1 Protein, His Tag (Cat. No. PD1-H522a) at 2 µg/mL (100 µL/well) can bind Human PD-L2, Mouse IgG1 Fc Tag (Cat. No. PD2-H52A5) with a linear range of 10-156 ng/mL (Routinely tested).


**Bioactivity-BLI**




Loaded Human PD-1 Protein, His Tag (Cat. No. PD1-H522a) on HIS1K Biosensor, can bind Human PD-L1, Fc Tag (Cat. No. PD1-H5258) with an affinity constant of 23.3 nM as determined in BLI assay (ForteBio Octet Red96e) (QC tested).



Loaded Human PD-1 Protein, His Tag (Cat. No. PD1-H522a) on HIS1K Biosensor, can bind Human PD-L2 Protein, Fc Tag (Cat. No. PD2-H5251) with an affinity constant of 15.7 nM as determined in BLI assay (ForteBio Octet Red96e) (QC tested).



Loaded Human PD-L1, Fc Tag (Cat. No. PD1-H5258) on Protein A Biosensor, can bind Human Human PD-1 Protein, His Tag (Cat. No. PD1-H522a) with an affinity constant of 4.8 µM as determined in BLI assay (ForteBio Octet Red96e) (Routinely tested).



Loaded Human PD-L2 Protein, Fc Tag (Cat. No. PD2-H5251) on Protein A Biosensor, can bind Human Human PD-1 Protein, His Tag (Cat. No. PD1-H522a) with an affinity constant of  $0.9 \mu\text{M}$  as determined in BLI assay (ForteBio Octet Red96e) (Routinely tested).

## Background

Programmed cell death protein 1 (PD-1) is also known as CD279 and PDCD1, is a type I membrane protein and is a member of the extended CD28/CTLA-4 family of T cell regulators. PDCD1 is expressed on the surface of activated T cells, B cells, macrophages, myeloid cells and a subset of thymocytes. PD-1 has two ligands, PD-L1 and PD-L2, which are members of the B7 family. PD-L1 is expressed on almost all murine tumor cell lines, including PA1 myeloma, P815 mastocytoma, and B16 melanoma upon treatment with IFN- $\gamma$ . PD-L2 expression is more restricted and is expressed mainly by DCs and a few tumor lines. PD1 inhibits the T-cell proliferation and production of related cytokines including IL-1, IL-4, IL-10 and IFN- $\gamma$  by suppressing the activation and transduction of PI3K/AKT pathway. In addition, coligation of PD1 inhibits BCR-mediated signal by dephosphorylating key signal transducer. In vitro, treatment of anti-CD3 stimulated T cells with PD-L1-Ig results in reduced T cell proliferation and IFN- $\gamma$  secretion. Monoclonal antibodies targeting PD-1 that boost the immune system are being developed for the treatment of cancer.

## Clinical and Translational Updates

Please contact us via [TechSupport@acrobiosystems.com](mailto:TechSupport@acrobiosystems.com) if you have any question on this product.