



Synonym

CDw328,D-siglec,A79 membrane protein,p75,Adhesion inhibitory receptor molecule 1, AIRM-1

Source

Human Siglec-7, His Tag(SG7-H5225) is expressed from human 293 cells (HEK293). It contains AA Gln 19 - Leu 353 (Accession # [Q9Y286-1](#)).
Predicted N-terminus: Gln 19

Molecular Characterization

Siglec-7(Gln 19 – Leu 353)
Q9Y286-1 Poly-his

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 38.8 kDa. The protein migrates as 55-65 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>95% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in 25 mM MES, 150 mM NaCl, pH5.5 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.


SDS-PAGE



Human Siglec-7, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

Bioactivity-ELISA

SEC-MALS




The purity of Human Siglec-7, His Tag (Cat. No. SG7-H5225) is more than 85% and the molecular weight of this protein is around 90-110 kDa verified by SEC-MALS.

[Report](#)

Discounts, Gifts,
and more!





Immobilized Human Siglec-7, His Tag (Cat. No. SG7-H5225) at 5 µg/mL (100 µL/well) can bind Neu5Aca2-6GalNAca-PAA-biotin with a linear range of 6-100 ng/mL (QC tested).

Background

Siglec-7 is a member of the human CD33-related Siglec receptor. The extracellular region of Siglec-7 is characterized by an N-terminal V-set Ig domain that can bind sialic acid and two C2-set Ig domains. The cytoplasmic tail of Siglec-7 has one immune-receptor tyrosine-based inhibitory motif (ITIM) and one ITIM-like motif. Siglec-7 is considered as a sialic acid-dependent immunoreceptor with inhibitory potential and expressed predominantly on human NK cells, monocytes and a small subset of CD8⁺ T cells.

Clinical and Translational Updates

Discounts, Gifts,
and more!

