

Synonym

Spike, Sprotein, Spike glycoprotein, Sglycoprotein

Source

SARS-CoV-2 Spike Trimer, His Tag (B.1.1.529/Omicron) (SPN-C52Hz) is the ectodomain of SARS-CoV-2 spike protein which contains AA Val 16 - Pro 1213 (Accession # QHD43416.1 (A67V, HV69-70del, T95I, G142D, VYY143-145del, N211del, L212I, ins214EPE, G339D, S371L, S373P, S375F, K417N, N440K, G446S, S477N, T478K, E484A, Q493R, G496S, Q498R, N501Y, Y505H, T547K, D614G, H655Y, N679K, P681H, N764K, D796Y, N856K, Q954H, N969K, L981F)). The 35 spike mutations are identified on the SARS-CoV-2 Omicron variant (Pango lineage: B.1.1.529; GISAID clade: GR/484A; Nextstrain clade: 21K). The recombinant protein is expressed from human 293 cells (HEK293) with T4 fibritin trimerization motif and a polyhistidine tag at the C-terminus. Proline substitutions (F817P, A892P, A899P, A942P, K986P, V987P) and alanine substitutions (R683A and R685A) are introduced to stabilize the trimeric prefusion state of SARS-CoV-2 S protein and abolish the furin cleavage site, respectively.

Molecular Characterization

Predicted N-terminus: Val 16

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 138.2 kDa. The protein migrates as 160-190 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>90% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.


SDS-PAGE



SARS-CoV-2 Spike Trimer, His Tag (B.1.1.529/Omicron) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%.

Bioactivity-ELISA


SEC-MALS




The purity of SARS-CoV-2 Spike Trimer, His Tag (B.1.1.529/Omicron) (Cat. No. SPN-C52Hz) is more than 90% and the molecular weight of this protein is around 525-605 kDa verified by SEC-MALS.

Report







Immobilized SARS-CoV-2 Spike Trimer, His Tag (B.1.1.529/Omicron) (Cat. No. SPN-C52Hz) at 1 μ g/mL (100 μ L/well) can bind Human ACE2, Fc Tag (Cat. No. AC2-H5257) with a linear range of 0.4-50 ng/mL (QC tested).



Immobilized SARS-CoV-2 Spike Trimer, His Tag (B.1.1.529/Omicron) (Cat. No. SPN-C52Hz) at 1 μ g/mL (100 μ L/well) can bind Anti-SARS-CoV-2 Spike RBD Antibody, Mouse IgG1 (Omicron Specific) (Cat. No. SPD-M305) with a linear range of 0.5-8 ng/mL (Routinely tested).





Anti-SARS-CoV-2 Spike RBD Antibody, Chimeric mAb, Human IgG1 Conc. (ng/mL)

Immobilized SARS-CoV-2 Spike Trimer, His Tag (B.1.1.529/Omicron) (Cat. No. SPN-C52Hz) at 1 μ g/mL (100 μ L/well) can bind Anti-SARS-CoV-2 Spike RBD Antibody, Chimeric mAb, Human IgG1 (Cat. No. S1N-M122) with a linear range of 0.4-50 ng/mL (Routinely tested).

Background

It's been reported that SARS-CoV-2 can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity.

Clinical and Translational Updates

