



Synonym

Streptavidin,SA

Source

Streptavidin Protein-Texas Red(STN-NT113) is expressed from E. coli cells.

Molecular Characterization

This protein carries no "tag".

The protein has a calculated MW of 6.72.

Conjugate

Texas Red

Excitation Wavelength: 586 nm

Emission Wavelength: 603 nm

Labeling

The primary amines in the side chains of lysine residues and the N-terminus of the protein are conjugated with Texas Red using standard chemical labeling method. The residual Texas Red is removed by molecular sieve treatment during purification process.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>95% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please protect from light and avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

Bioactivity-FACS



5e5 of Anti-CD19 CAR-293 cells were stained with 100 µL of 20 µg/mL Biotinylated Human CD19 (20-291) Protein, Fc, Avitag™, premium grade (Cat. No. CD9-H82F6) and negative control protein respectively, washed and then followed with 1 µg/mL of Streptavidin Protein-Texas Red (Cat. No. STN-NT113) and analyzed with FACS. Texas Red signal was used to evaluate the binding activity (QC tested).

Discounts, Gifts,
and more!





Background

Streptavidin is a 66KDa tetrameric protein purified from the bacterium *Streptomyces avidinii*, and exhibits high binding affinity to biotin. Each unit can bind one biotin. Horseradish peroxidase is metalloenzyme, a 44KDa glycoprotein. When incubate with substrates, it produces a coloured, fluorimetric, or luminescent derivatives, which can be detected and quantified. HRP conjugated Streptavidin is widely used for the detection and quantification of biotinylated proteins.

Clinical and Translational Updates

Please contact us via TechSupport@acrobiosystems.com if you have any question on this product.

Discounts, Gifts,
and more!

