

**Synonym**

CEDLAP,TGF-beta 1,TGFB1,DPD1,TGF-beta-1,TGFB

Source

Human TGF-Beta 1 Protein, Tag Free(TG1-H4212) is expressed from human 293 cells (HEK293). It contains AA Ala 279 - Ser 390 (Accession # [P01137-1](#)).
 Predicted N-terminus: Ala 279

Molecular Characterization

TGFB1(Ala 279 - Ser 390)
P01137-1

This protein carries no "tag".

The protein has a calculated MW of 12.8 kDa (monomer). The protein migrates as 14 kDa (monomer) under reducing (R) condition, and 25 kDa (Dimer) when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under non-reducing (NR) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 0.1 EU per µg by the LAL method.

Host Cell Protein

<0.5 ng/µg of protein tested by ELISA.

Host Cell DNA

<0.02 ng/µg of protein tested by qPCR.

Sterility

Negative

Mycoplasma

Negative.

Purity

>95% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in 0.085% TFA in 30% ACN with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.


Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 12 months under sterile conditions after reconstitution.


SDS-PAGE

Human TGF-Beta 1 Protein, Tag Free on SDS-PAGE under reducing (R) and non-reducing (NR) conditions. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95% (With [Star Ribbon Pre-stained Protein Marker](#)).

Bioactivity-ELISA


Discounts, Gifts,
and more!





Immobilized Human TGF-Beta 1 Protein, Tag Free (Cat. No. TG1-H4212) at 0.2 µg/mL (100 µL/well) can bind Human TGFBR2, Fc Tag (Cat. No. TG2-H5252) with a linear range of 0.2-2.5 ng/mL (QC tested).


Bioactivity-SPR



Human TGF-Beta 1 Protein, Tag Free (Cat. No. TG1-H4212) immobilized on CM5 Chip can bind Human TGF-beta RI Protein, Fc Tag (Cat. No. TG1-H5254) with an affinity constant of 99.2 nM as determined in a SPR assay (Biacore 8K) (Routinely tested).

Bioactivity-Bioactivity CELL BASE

Human TGFB1 Protein inhibits the IL-4 dependent proliferation of TF-1 cells



Human TGF-Beta 1 Protein, Tag Free (Cat. No. TG1-H4212) inhibits the Human IL-4, premium grade (Cat. No. IL4-H4218) dependent proliferation the TF-1 cells. The specific activity of Human TGF-Beta 1 Protein, Tag Free is >

Discounts, Gifts,
and more!



Human TGF-Beta 1 / TGFB1 Protein, Tag Free

Catalog # TG1-H4212



BIOSYSTEMS
Acro

8.00x10⁶ IU/mg, which is calibrated against transforming growth factor β 1 (NIBSC code: 89/514) (QC tested).

Background

Transforming growth factor beta 1 (TGFB1) is also known as TGF- β 1, CED, DPD1, TGFB. is a polypeptide member of the transforming growth factor beta superfamily of cytokines. It is a secreted protein that performs many cellular functions, including the control of cell growth, cell proliferation, cell differentiation and apoptosis. The TGFB1 protein helps control the growth and division (proliferation) of cells, the process by which cells mature to carry out specific functions (differentiation), cell movement (motility), and the self-destruction of cells (apoptosis). The TGFB1 protein is found throughout the body and plays a role in development before birth, the formation of blood vessels, the regulation of muscle tissue and body fat development, wound healing, and immune system function. TGFB1 is particularly abundant in tissues that make up the skeleton, where it helps regulate bone growth, and in the intricate lattice that forms in the spaces between cells (the extracellular matrix). Within cells, this protein is turned off (inactive) until it receives a chemical signal to become active. TGFB1 plays an important role in controlling the immune system, and shows different activities on different types of cell, or cells at different developmental stages. Most immune cells (or leukocytes) secrete TGFB1. TGFB1 has been shown to interact with TGF beta receptor 1, LTBP1, YWHAE, EIF3I and Decorin.

Clinical and Translational Updates

Discounts, Gifts,
and more!



www.acrobiosystems.com

8/8/2024