

Synonym

DIF, TNF-alpha, TNFA, TNFSF2, cachexin, cachectin, TNF α

Source

Human TNF-alpha Protein, Fc Tag(TNA-H5255) is expressed from human 293 cells (HEK293). It contains AA Val 77 - Leu 233 (Accession # [P01375-1](#)).

Predicted N-terminus: Val 77

Molecular Characterization

TNF-alpha(Val 77 - Leu 233) P01375-1	Fc(Pro 100 - Lys 330) P01857
---	---------------------------------

This protein carries a human IgG1 Fc tag at the C-terminus

The protein has a calculated MW of 43.8 kDa. The protein migrates as 50-53 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per μ g by the LAL method.

Purity

>90% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 μ m filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- 20°C to -70°C for 12 months in lyophilized state;
- 70°C for 3 months under sterile conditions after reconstitution.


SDS-PAGE



Human TNF-alpha Protein, Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90% (With [Star Ribbon Pre-stained Protein Marker](#)).


Bioactivity-ELISA

SEC-MALS



The purity of Human TNF-alpha Protein, Fc Tag (Cat. No. TNA-H5255) is more than 90% and the molecular weight of this protein is around 255-285 kDa verified by SEC-MALS.

[Report](#)



Immobilized Human TNFR1, His Tag (Cat. No. TN1-H5222) at 5 µg/mL (100 µL/well) can bind Human TNF-alpha Protein, Fc Tag (Cat. No. TNA-H5255) with a linear range of 0.1-2 ng/mL (QC tested).

Background

Tumor necrosis factor alpha (TNF α) is a cytokine produced primarily by monocytes and macrophages. It is found in synovial cells and macrophages in the tissues. The primary role of TNF α is in the regulation of immune cells. TNF α is able to induce apoptotic cell death, to induce inflammation, and to inhibit tumorigenesis and viral replication. Dysregulation of TNF α production has been implicated in a variety of human diseases, including major depression, Alzheimers disease and cancer. Recombinant TNF α is used as an immunostimulant under the INN tasonermin. TNF α can be produced ectopically in the setting of malignancy and parallels parathyroid hormone both in causing secondary hypercalcemia and in the cancers with which excessive production is associated.

Clinical and Translational Updates

Please contact us via TechSupport@acrobiosystems.com if you have any question on this product.